

Dynamics of particles and manifolds in a quenched random force field

Pierre Le Doussal[†], Leticia F. Cugliandolo[‡], and Luca Peliti[¶]

[†] CNRS-Laboratoire de Physique Théorique de l'Ecole
Normale Supérieure, 24 rue Lhomond, F-75231 Paris*

[‡] Laboratoire de Physique Théorique des Liquides, Jussieu
4, Place Jussieu, 75005 Paris France**

[¶] Groupe de Physico-Chimie Théorique, CNRS URA 1382, ESPCI, 10, rue Vauquelin, F-75231 Paris Cedex 05, France,***
and

Dipartimento di Scienze Fisiche, Unità INFN, Università "Federico II", Mostra d'Oltremare, Pad. 19, I-80125 Napoli, Italy.
(February 1, 2008)

We study the dynamics of a directed manifold of internal dimension D in a d -dimensional random force field. We obtain an exact solution for $d \rightarrow \infty$ and a Hartree approximation for finite d . They yield a Flory-like roughness exponent ζ and a non trivial anomalous diffusion exponent ν continuously dependent on the ratio g_T/g_L of divergence-free (g_T) to potential (g_L) disorder strength. For the particle ($D = 0$) our results agree with previous order ϵ^2 RG calculations. The time-translational invariant dynamics for $g_T > 0$ smoothly crosses over to the previously studied ultrametric aging solution in the potential case.

74.60.Ge, 05.20.-y

The problem of the diffusion of a particle in a d -dimensional static random force field has been often addressed in the past decade [1–6]. It is of interest in problems like tracer diffusion in porous media [1,7] or turbulent flows [8]. Typically the force is the sum of a potential random force (longitudinal in k -space) of strength g_L and a divergence-free (transverse) one of strength g_T . The diffusion is anomalous [1–6] ($r \sim t^\nu$ with $\nu \neq 1/2$) if either (i) the dimension d of space is low ($d \leq 2$), or (ii) the force field has long-range correlations ($\overline{F(0)F(r)} \sim r^{-a}$ with $a \leq 2$). While this is true for weak gaussian disorder, other anomalous behaviors are possible for disorder distributions with tails [5]. Apart from particular cases (such as the long-range correlated potential case [4], including $d = 1$ [6]), most of the above results were obtained by perturbation theory in the disorder around the thermal value $\nu_{th} = 1/2$, which holds in the quadrant $d > 2$ and $a > 2$. An expansion in powers of $\epsilon' = 2 - d$ was performed for short range (SR) disorder [1], and one in powers of $\epsilon = 2 - a$ for long range (LR) correlated disorder [2,3] (and a double ϵ, ϵ' expansion [3] near the point $d = 2, a = 2$). Superdiffusive behavior was obtained for $g_T/g_L \gg 1$ and subdiffusive slow dynamics for $g_T/g_L \ll 1$. A line of fixed points was obtained up to order ϵ^2 , with a continuously variable exponent ν , in the case of LR correlations ($d > 2, a < 2$), parametrized by the ratio g_T/g_L . The divergence-free case $g_L = 0$ is simpler, since the value of ν is equal to the Flory value ν_F obtained by purely dimensional arguments, as discussed below.

On the other hand, out of equilibrium relaxational (potential) dynamics has recently been studied extensively, in the context of spin glass models [9] as well as particles [10,11] and manifolds [12] in random potentials. Analytical solutions were obtained [9] for the long-time dynamics in mean-field models, showing explicitly their glassy features such as slow (aging) dynamics, and the breakdown of time-translation invariance (TTI) and of fluctuation-dissipation theorems (FDT). Whether these glassy properties of potential dynamics survive to a small *non-potential* perturbation was recently addressed [13], and it was found that, while the dynamics becomes TTI in most cases, some glassy features survive. Similarly, a glassy state with stationary dissipation was proposed [14] for driven vortex systems.

In this paper we thus reconsider the dynamics of a particle ($D = 0$) in a d -dimensional random force field as well as its extension to a directed elastic manifold of internal dimension D . We obtain the diffusion exponent $r \sim t^\nu$ and (for $D > 0$) the roughness exponent $r \sim x^\zeta$ where x is the internal coordinate. Our results are exact for $d \rightarrow \infty$. This mean-field limit is non trivial if transverse and longitudinal parts are scaled properly. It extends the previous results [2,3] obtained via RG since it allows to reach *nonperturbatively* arbitrary values of the ratio g_T/g_L (that the RG could not reach because of runaway flows for $g_T = 0$). We can now explore systematically the interesting potential limit $g_T \rightarrow 0$ and we prove the remarkable property of continuity of the mean-field dynamic solution when aging is approached for $g_T \rightarrow 0$. This property was proposed in [13] and verified in numerical solutions of the p -spin model. We suggest to take systematically advantage of it, in order to improve numerical simulations by adding a small nonpotential part, waiting till TTI is attained, and taking the nonpotential part to zero *after* the large size limit. We find that the finite- d Hartree approximation compares well with the RG.

We study the Langevin dynamics of a directed elastic manifold, of internal dimension D , parametrized by a d -component field $r(x)$:

$$\partial_t r_\alpha(x, t) = \nabla_x^2 r_\alpha(x, t) + F_\alpha[r(x, t), x] + \eta_\alpha(x, t) , \quad (1)$$

where η is the gaussian thermal noise, satisfying $\langle \eta_\alpha(x, t) \eta_\beta(x', t') \rangle = 2T \delta_{\alpha\beta} \delta(t - t') \delta^D(x - x')$, and $F_\alpha[r, x]$ is the gaussian quenched random force with a correlator $\overline{F_\alpha[r, x] F_\beta[r', x']} = \Delta_{\alpha\beta}(r - r') \delta^D(x - x')$ which depends only on the distance $|r - r'|$. We denote by angular brackets averages over the thermal noise and by an overbar averages over the random force. The force has both a potential (L) and a divergence-free (T) part, which contribute separately to the correlator:

$$\Delta_{\alpha\beta}(r) = -\partial_{r_\alpha} \partial_{r_\beta} \Delta_L(r) - \left(\delta_{\alpha\beta} \sum_\gamma \partial_{r_\gamma}^2 - \partial_{r_\alpha} \partial_{r_\beta} \right) \Delta_T(r). \quad (2)$$

We study the case in which the correlations at large scale behave as power laws: the Fourier transforms of the correlators with respect to r behave, for small values of their argument K , as $\Delta_{\alpha\beta}(K) \sim K^{a-d} (\tilde{g}_L P_{\alpha\beta}^L(K) + \tilde{g}_T P_{\alpha\beta}^T(K))$, where $P_{\alpha\beta}^L(K) = K_\alpha K_\beta / K^2$ is the longitudinal projector and $P_{\alpha\beta}^T(K) = 1 - P_{\alpha\beta}^L(K)$ is the transverse one. We compute the mean-squared displacement $\overline{D_{xx'}(t, t')}$ and the correlation $\overline{B_{xx'}(t, t')}$, respectively defined by $D_{xx'}(t, t') = \frac{1}{d} \langle (r(x, t) - r(x', t'))^2 \rangle$ and $B_{xx'}(t, t') = \frac{1}{d} \langle (r(x, t) - r(x', t')) \cdot (r(x', t) - r(x', t')) \rangle$. The mean-square *displacement* at equal *times* $D_{xx'}(t, t) \sim |x - x'|^{2\zeta}$ measures the roughness of the manifold, while the *correlation* at equal *space* $\tilde{B}(t, t') = B_{xx}(t, t')$ measures the diffusion of a single monomer. When TTI holds, the diffusion exponent ν is defined by $\tilde{B}(t, t') \sim |t - t'|^{2\nu}$. In the absence of disorder one has $\zeta = \zeta_{\text{th}} = (2 - D)/2$, $\nu = \nu_{\text{th}} = (2 - D)/4$ for $D < 2$, and bounded motion and fluctuations for $D > 2$. Before proceeding, let us consider the problem by Flory-like dimensional arguments. If all terms in (1) scaled the same way one would have $r/t \sim r/x^2 \sim r^{-\gamma} x^{-D/2}$ with $\gamma = a/2$ for $d > a$. This yields

$$\zeta_F = 2\nu_F = \frac{4 - D}{4(1 + \gamma)}. \quad (3)$$

It turns out that, for $d \rightarrow \infty$, $\zeta = \zeta_F$ is exact but $\nu = \nu_F$ holds only in the transverse case.

One can take the large- d limit directly in the De Dominicis-Janssen functional, performing a gaussian decoupling of $r(x, t)$ and the conjugate field $\hat{r}(x, t)$. We define:

$$\Delta_T(r) = -V_T(r^2/d), \quad \Delta_L(r) = -d V_L(r^2/d). \quad (4)$$

One can also study in any d the self-consistent Hartree approximation where $r(x, t)$ and $\hat{r}(x, t)$ are treated as Gaussian variables. It amounts to replace everywhere, in the equations and solutions obtained below for $d = \infty$, $V_{L,T}$ by $\tilde{V}_{L,T}$ defined by $\tilde{V}(z) = \int d^d \phi / (2\pi z)^{d/2} \exp(-\phi^2 / (2z)) V(\phi^2/d)$ (see, e.g., appendix A of [11]). The coupled dynamical equations also involve the response $R_{xx'}^{\alpha\beta}(t, t') = 1/d \overline{\delta \langle r^\alpha(x, t) \rangle / \delta f^\beta(x', t') |}_{f=0}$ where $f(x', t')$ is a small perturbation applied at point x' at time t' . The response and correlations are isotropic, $R_{xx'}^{\alpha\beta}(t, t') = (\delta_{\alpha\beta}/d) R_{xx'}(t, t')$. We choose a space-homogeneous initial condition $B_{xx'}(0, 0) = B_{x-x'}(0, 0)$ and denote by B_k and R_k the Fourier transforms with respect to $x - x'$, and by tilde equal space ($x = x'$) two-time functions (e.g., $\tilde{B} = \int_k B_k$). We use the shorthand $\int_k \equiv \int d^D k / (2\pi)^D$ and $\int_\omega \equiv \int d\omega / (2\pi)$. With $R_k(t, t) = 0$ and $R_k(t, t^-) = 1$, the dynamical equations read

$$\partial_t R_k(t, t') = -k^2 R_k(t, t') - \int_0^t ds \Sigma(t, s) (R_k(t, t') - R_k(s, t')) + \delta(t - t'), \quad (5)$$

$$\begin{aligned} \partial_t B_k(t, t') = & -k^2 B_k(t, t') + 2T + \int_0^t ds D(t, s) R_k(t, s) - \int_0^{t'} ds D(t, s) R_k(t', s) \\ & - \int_0^t ds \Sigma(t, s) (B_k(t, s) - B_k(s, t') + B_k(t, t')) - 4T R_k(t', t), \end{aligned} \quad (6)$$

where $D(t, s) = 4V_1'(\tilde{B}(t, s))$ and $\Sigma(t, s) = -4V_2''(\tilde{B}(t, s))\tilde{R}(t, s)$. We define $V_1 = V_L + (1 - 1/d)V_T$ and $V_2 = V_L$, and consider $V_1(b) = g_1 b^{1-\gamma}/(2(1-\gamma))$, $V_2(b) = g_2 b^{1-\gamma}/(2(1-\gamma))$ where $g_1 = g_L + (1 - 1/d)g_T$, $g_2 = g_L$, and $\gamma = a/2$ for large d , thus generalizing the potential case studied in [11,12]. We define $\tilde{g}_{L,T} = -g_{L,T} C_{d,\gamma}/(2(1-\gamma))$, where $C_{d,\gamma} = \int_r (r^2/d)^{1-\gamma} \exp(in \cdot r)$ with $n^2 = 1$. As shown below, the case of LR correlations corresponds (for $d \rightarrow \infty$) to $\gamma > 2/(2 - D)$ for $D < 2$, and any γ for $D > 2$.

When $g_T \neq 0$ one can assume that a TTI solution is reached at large t' , $B_k(t, t') = B_k(t - t')$ and $R_k(t, t') = R_k(t - t')$. We checked this property numerically for $D = 0$. However we stress that it is not certain that it should hold for any nonpotential dynamics [16]. Using TTI one can recast the above equations in the form

$$R_k(\omega) = \frac{1}{-i\omega + k^2 + \Sigma(0) - \Sigma(\omega)}, \quad B_k(\omega) = \frac{2T + D_k}{k^2} 2\pi\delta(\omega) - \frac{4T + D(\omega)}{|-i\omega + k^2 + \Sigma(0) - \Sigma(\omega)|^2},$$

$$\Sigma(\omega) = -4 \int_{-\infty}^{+\infty} dt \exp(i\omega t) V_2''(b(t)) r(t), \quad D(\omega) = 4 \int_{-\infty}^{+\infty} dt \exp(i\omega t) V_1'(b(t)), \quad (7)$$

where $D_k = \int_{\omega'} (D(\omega') R_k(-\omega') - \Sigma(\omega') B_k(\omega'))$. The correlation $C_k(\omega) = 1/d \langle \overline{(r(k, \omega) \cdot r(-k, -\omega))}$ thus obeys $2C_k(\omega) = (4T + D(\omega))/|-i\omega + k^2 + \Sigma(0) - \Sigma(\omega)|^2$. The displacement can then be computed using $D_x(t) = 2 \int_{k, \omega} (1 - \cos(kx + \omega t)) C_k(\omega)$. In the large time limit we propose

$$\tilde{R}(t) \sim R t^{-\mu}, \quad \tilde{B}(t) \sim B t^{2\nu}, \quad (8)$$

and define the FDT violation factor $X[B]$ by $\tilde{R}(t) = X[\tilde{B}(t)] \partial_t \tilde{B}(t)$. Equation (8) implies the small frequency behaviors $\tilde{R}(\omega) \sim R \Gamma[1 - \mu] (-i\omega)^{\mu-1}$ for $\mu < 1$ or $\tilde{R}(\omega) - \tilde{R}(0) \sim R \Gamma[1 - \mu] (-i\omega)^{\mu-1}$ for $\mu > 1$ and $\tilde{B}(\omega) \sim 2B \Gamma[1 + 2\nu] \text{Re}(i\omega)^{-1-2\nu}$. It also implies $D(\omega) \sim 8g_1 B^{-\gamma} \Gamma[1 - 2\nu\gamma] \text{Re}(i\omega)^{2\nu\gamma-1}$ and $\Sigma(0) - \Sigma(\omega) \sim -4g_2 \gamma R B^{-(1+\gamma)} \Gamma[1 - \mu - 2\nu(\gamma + 1)] (-i\omega)^{\mu-1+2\nu(\gamma+1)}$. One must have $0 < \mu - 1 + 2\nu(\gamma + 1) < 1$ in order to be able to neglect the $-i\omega$ in the Green functions and to have a finite $\Sigma(0)$. We shall find out that this is always the case for $g_2 > 0$. The equation for $\tilde{R} = \int_k R_k$ yields, at small ω and for $D < 2$,

$$\tilde{R}(\omega) \sim A_D (\Sigma(0) - \Sigma(\omega))^{\frac{D-2}{2}}, \quad (9)$$

with $A_D = \pi/(2^D \pi^{D/2} \Gamma[D/2] \sin(\pi D/2))$. For $D > 2$, $\tilde{R}(0)$ is finite and one has instead $\tilde{R}(\omega) - \tilde{R}(0) \sim A_D (\Sigma(0) - \Sigma(\omega))^{\frac{D-2}{2}}$. This gives the relation between exponents:

$$1 - \mu = 2\nu(\gamma + 1) \frac{2 - D}{4 - D}. \quad (10)$$

This value for μ is consistent with neglecting the $-i\omega$ term in (7), provided that $0 < \nu < \nu_T$, where $\nu_T = (4 - D)/(4(1 + \gamma))$. It implies $X[\tilde{B}] \sim X\tilde{B}^w$ with $X = R/(2\nu) B^{-(1+\gamma)\frac{2-D}{4-D}}$ and $w = -1 + (1 + \gamma)\frac{2-D}{4-D}$. It also yields the following relation between amplitudes:

$$(2\nu X)^{\frac{4-D}{2}} = A_D \frac{(-4g_2 \gamma \Gamma[2\nu(\gamma + 1)\frac{2-D}{4-D}])^{\frac{D-2}{2}}}{\Gamma[2\nu(\gamma + 1)\frac{2-D}{4-D}]}. \quad (11)$$

Similarly the equation for $\tilde{B} = \int_k B_k$ yields

$$\tilde{B}(\omega) \sim A_D D(\omega) |\Sigma(0) - \Sigma(\omega)|^{\frac{D-4}{2}} \frac{\sin(\phi \frac{2-D}{2})}{\sin(\phi)}, \quad (12)$$

where $\phi = -\frac{\pi}{2}(\mu - 1 + 2\nu(\gamma + 1))$ is the phase of $\Sigma(0) - \Sigma(\omega)$. One can check that the exponent relation (10) is now automatically satisfied. The resulting relation between amplitudes, together with the relation (11), yields our main result, i.e., the equation which determines the exponent ν :

$$\frac{g_2}{g_1} = h(\nu, D, \gamma) \equiv \frac{\Gamma[2\nu(\gamma + 1)\frac{2-D}{4-D}]}{|\Gamma[2\nu(\gamma + 1)\frac{2-D}{4-D}]|} \frac{\Gamma[1 - 2\nu\gamma] \sin(\pi\gamma\nu)}{\Gamma[1 + 2\nu\gamma] \sin(\pi\nu)} \frac{\sin(\frac{\pi}{2} 2\nu(1 + \gamma)\frac{2-D}{4-D})}{\sin(\frac{\pi}{2} 2\nu(1 + \gamma)\frac{2-D}{4-D})}, \quad (13)$$

where, we recall, $g_2/g_1 = g_L/(g_L + g_T)$ when $d \rightarrow \infty$. The function h is a monotonic function of ν , such that $h(\nu = 0, D, \gamma) = 1$ and $h(\nu = \nu_F, D, \gamma) = 0$ where ν_F is given by (3). Thus we find that in the transverse case $\nu = \nu_T \equiv \nu_F$ and that ν spans the interval $0 < \nu < \nu_T$ when the ratio g_2/g_1 varies from 1 (potential case) to 0 (divergence free case). These results hold in the LR case $\gamma < \gamma_c$. The limit $\gamma \rightarrow \gamma_c^-$ (for which $h(\nu, D, \gamma_c) = 1$) is studied below. One also easily obtains the roughness exponent ζ . From $C_k(\omega)$ and Eq. (7) one finds that $t \sim x^z$ with $z = 2\nu/\nu_T$ and $r \sim t^\nu$. Thus $\zeta = z\nu = 2\nu_T = \zeta_F$. This is the Flory value, as announced, and interestingly, is found not to depend on the precise nature of the random force (though the physics is quite different as the ratio g_L/g_T varies). For finite d our approach also yields a Hartree approximation which amounts to: (i) replace in (13) g_1 by $g_L + (1 - \frac{1}{d})g_T$ and g_2 by g_L ; (ii) setting $\gamma = \min(a, d)/2$ ($\gamma = d/2$ corresponds to the Hartree SR case and $\gamma < d/2$ the Hartree LR case since the model studied here and in [2,3] is a random *force* model).

Let us compare our result for ν (exact when $d \rightarrow \infty$) with the RG calculations for the particle ($D = 0$) with $\gamma = 1 - \epsilon$ in any d . The RG calculation was carried out upto order ϵ in [2] and upto order ϵ^2 in [3]. In our notation, it reads [3]

$$\frac{1}{2\nu} = 1 - \frac{d-1-dk}{2(d-1)}\epsilon + \epsilon^2 dk \frac{4+2d-5d^2+d^3+dk(4-2d+d^2)}{4(2-d)(d-1)^2d}, \quad (14)$$

where $k = g_L/g_T$, while, from (13) and using the above Hartree values one finds

$$\frac{1}{2\nu} = 1 - \frac{g_1-2g_2}{2(g_1-g_2)}\epsilon - \frac{g_2g_1\epsilon^2}{4(g_1-g_2)^2} = 1 - \frac{d-1-dk}{2(d-1)}\epsilon + \frac{\epsilon^2 dk(1-d-dk)}{4(d-1)^2}. \quad (15)$$

For $d \rightarrow \infty$, ν from (13) and (14) coincide. ν is determined by the bare ratio g_T/g_L , which indeed is unrenormalized in the RG. Furthermore the finite d Hartree approximation is found to be *exact to first order* in $\epsilon = 1 - \gamma$ for all $d > 2$. Thus it is a good approximation in the Hartree LR case and is exact for $d = 1$ since it yields $\nu = 0$. Though it does predict correctly the special point $d = 2, \gamma = 1$ it is not able to reproduce correctly the delicate crossover [3] around this point. The RG approach of [3,2] can be generalized to the manifold, and for $0 < D < 2$ one can similarly expand around $\gamma_c(D) = 2/(2-D)$ (LR) (and around $d_c(D) = 4/(2-D)$ (SR)). The result (13) leads to the following formula, exact for $d \rightarrow \infty$, with $\gamma = 2/(2-D) - \epsilon$ to first order in ϵ :

$$\nu = \frac{2-D}{4} - \epsilon \frac{(2-D)^2}{8(4-D)} \frac{(4-D)g_2-2g_1}{g_1-g_2}. \quad (16)$$

It would be interesting to check whether the Hartree approximation is also exact for any $D < 2$, d to $O(\epsilon)$.

Up to now we have studied the limit $\gamma \rightarrow \gamma_c(D)$ keeping the ratio $g_L/g_T > 0$ fixed, in which case $\nu \rightarrow \nu_{th}$. In fact the limit $\gamma \rightarrow \gamma_c$ is very non uniform, and therefore much richer. For $D < 2$ one finds that any value of ν with $0 < \nu < \nu_{th}$ can be reached if one takes the limit $\gamma \rightarrow \gamma_c(D)$ with the ratio $g_T/\epsilon g_L$ fixed. For the particle, this is perfectly consistent with the RG result obtained [3,2] in any d , which in our notations read $1/\nu = 1 + g_L + O(g_L^3)$ (note that it has a nice $d \rightarrow \infty$ limit and that there is a conjecture [3,2] that all higher order terms vanish). For $2 < D < 4$ one finds that the limit $\gamma \rightarrow \infty$ is non trivial with $\nu\gamma$ a function of g_T/g_L . Finally the limit $D = 4 - \epsilon$ is also very non uniform and any value of ν can be reached if g_T/ϵ is kept fixed as $\epsilon \rightarrow 0$.

We now study the potential limit $g_T \rightarrow 0$ and show on the case $D = 0$ that one goes continuously to the aging dynamics which holds for $g_T = 0$. This is best illustrated using the formalism of Ref. [12]. Both in the case of a solution satisfying ($g_T > 0$) and in the asymptotic aging solution ($g_T = 0$), one defines the FDT violation factor $X[B]$ and the triangular relation $\bar{f}(B, B')$ by

$$X[B] = \lim_{\substack{t \geq t' \rightarrow \infty \\ B(t, t') = B}} R(t, t') / \partial_{t'} B(t, t'), \quad \bar{f}(B', B) = \lim_{\substack{t \geq s \geq t' \rightarrow \infty \\ B(t, t') = B \geq B(t, s) = B'}} B(s, t'). \quad (17)$$

The dynamical equations are naturally written in terms of the linear susceptibility $\chi(t, t') = \int_{t'}^t ds R(t, s)$, which takes the form $\chi(B) = \int_0^B dB' X(B')$, and of the “anomaly” $M[B] = 4 \int_0^B dB' V''(B') X(B')$. The equation for R only contains V_2 and is thus identical to the case $g_T = 0$ and reads, in the long time limit, like Eq. (5.3) of Ref. [11]:

$$0 \approx \frac{d\chi[B]}{dt} = -1 - \chi[B] \int_B^\infty dB' 4V_2''(B') X[B'] + \int_0^B dB' 4V_2''(B') X[B'] (\chi[\bar{f}(B', B)] - \chi[B]). \quad (18)$$

We used $B(t, 0) \rightarrow \infty$. This equation is coupled to a similar (more cumbersome) equation for B that also involves V_1 . We specialize to the long range case $\gamma < 1$. The integrals are naively divergent at large B since $\lim_{B \rightarrow \infty} \chi[B] = \infty$ [15]. In the equation for B similar divergences occur. However, at large B , one finds $\chi[B] \sim 2X/(1+\gamma)B^{(1+\gamma)/2}$, as well as $\bar{f}(B', B) = (B^{\frac{1}{2\gamma}} - B'^{\frac{1}{2\gamma}})^{2\gamma}$. The divergent parts cancel and one recovers the equation for the amplitudes (11) (independent of small B details). This solution interestingly shows that when $\nu \rightarrow 0$ as $g_T \rightarrow 0$ the triangular relation continuously becomes $\bar{f}(B', B) = \max(B', B)$, i.e. the known ultrametric solution for the potential case. Similarly $X[B] \sim XB^{(\gamma-1)/2}$ tends continuously to the result of [11].

In the transverse case $g_L = 0$ for the particle $D = 0$ our method yields the Flory estimate $\nu = \nu_F \equiv 1/(1 + \min(a, d)/2)$, which turns out to be exact (for $\nu_F > 1/2$). This can either be seen perturbatively to all orders in the field theory [3], or, more physically, using that the fact that the *stationary* measure is exactly given by $P_{stat} = \text{const.}$ The Derrida-Luck method [17,18] then produces the result (and also yields rigorously the velocity-force law $v \sim F^\phi$

with $\phi = 1$) as we now show. One introduces a medium with random forces, but periodic with a large period Λ . An *exact* formula for the velocity in this medium is $v_\Lambda = \int dx F(x) P_{\text{stat}}$ where $P_{\text{stat}}(x)$ is a stationary measure (satisfying $0 = \nabla^2 P_{\text{stat}} - \partial_i (F_i P_{\text{stat}})$) with periodic boundary conditions. Using that $P_{\text{stat}} = 1/\Lambda^d$ one finds that the mean-squared velocity $\overline{v_\Lambda^2} \sim \Lambda^{-\min(a,d)}$ in that periodic medium (equivalently finite torus of volume Λ^d). This exact scaling relation ensures that $\nu = \nu_F$. For the manifold $D > 0$, however, the stationary measure is not known (the naive extension $P_{\text{stat}} \sim \exp(-\int dx (\nabla r)^2/2T)$ is not stationary). Thus the Flory values obtained here for the manifold in a divergence free flow may very well be exact only for $d \rightarrow \infty$.

One can also extend the arguments of Ref. [8] to the diffusion of an oriented (e.g. by an external field) manifold in a divergence-free turbulent flow. If the random force, proportional to the local velocity, can be considered as static on some time scales, with LR gaussian correlations $v(0)v(r) \sim r^{2/3+\mu/3}$ (μ arises from the intermittency corrections to Kolmogorov scaling), the result $\nu = \nu_T$ for the corresponding diffusion exponent yields $\nu = 3(4 - D)/(2(4 - \mu))$, which can be considered as a generalization of the Richardson's law (corrected by intermittency).

In conclusion, we have studied the problem of the dynamics of a manifold or a particle in a gaussian random force flow. We have obtained the roughness exponent ζ and ν exactly for $d \rightarrow \infty$. Our study establishes a bridge between several problems (aging and nonpotential dynamics) and techniques (large d and RG approaches).

We thank J. Kurchan for useful discussions. LFC thanks the SPHT-Saclay for hospitality and the European Union for financial support through the contract ERBCHRXCT920069. LP is Associato INFN, Sezione di Napoli, and acknowledges the support of a Chaire Joliot de l'ESPCI.

* Laboratoire Propre du CNRS, associé à l'Ecole Normale Supérieure et à l'Université Paris-Sud.

** Laboratoire Associé au CNRS. Also at SPEC, CEA, Saclay, France.

*** Laboratoire Associé au CNRS.

- [1] J.M. Luck; Nucl. Phys. **B225**, 169 (1983). J.A. Aronovitz and D.R. Nelson; Phys. Rev. **A30**, xx (1984). D.S. Fisher; Phys. Rev. **A30**, 960 (1984). V.E. Kratsov, I.V. Lerner and V. Yudson; J. Phys. **A18**, L703 (1985). L. Peliti; Phys. Rep. **103**, 225 (1984). D.S. Fisher, D. Friedan, Z. Qiu and S. Shenker; Phys. Rev. **A31**, 3841 (1985).
- [2] J.P. Bouchaud, A. Comtet, A. Georges and P. Le Doussal; J. Phys. (France) **48** 1445 (1987), *ibid* **49** 369 (1988).
- [3] J. Honkonen and E. Karjalainen; J. Phys. **A21**, 4217 (1988) and references therein.
- [4] W. Durett; Comm. Math. Phys. **104**, 87 (1986).
- [5] J. Machta; J. Phys. **A18**, L531 (1985).
- [6] J. P. Bouchaud, A. Comtet, A. Georges and P. Le Doussal; Ann Phys. **201**, 285 (1990).
- [7] S. Havlin and D. Ben Avraham; Adv. Phys. **36**, 695 (1987).
- [8] H. Hentschel and I. Procaccia; Phys. Rev. **A27**, 1266 (1983).
- [9] L.F. Cugliandolo and J. Kurchan; Phys. Rev. Lett. **71**, 173 (1993); J. Phys. **A27**, 5749 (1994).
- [10] S. Franz and M. Mézard; Physica **A209**, 1 (1994).
- [11] L. F. Cugliandolo and P. Le Doussal; Phys. Rev. **E53**, 1525 (1996).
- [12] L. F. Cugliandolo, J. Kurchan, P. Le Doussal; Phys. Rev. Lett. **76**, 2390 (1996) and to be published.
- [13] L. F. Cugliandolo, J. Kurchan, P. Le Doussal and L. Peliti; cond-mat/9606060, Phys. Rev. Lett. in press.
- [14] T. Giamarchi and P. Le Doussal; Phys. Rev. Lett. **76**, 3408 (1996).
- [15] This can be seen from Ref [11] by letting $\mu \rightarrow 0$ since with a mass μ one has $\chi[B[t, 0]] = 1/\mu$.
- [16] See for instance the $p = 2$ spherical spin model at $T = 0$ in [13].
- [17] B. Derrida and J.M. Luck; Phys. Rev. **B28**, 7183 (1983).
- [18] See applications of Ref. [17] in B. Derrida and Y. Pomeau, Phys. Rev. Lett. **48**, 627 (1982); P. Le Doussal and J. Machta Phys. Rev. **B40**, 9427 (1989); P. Le Doussal and V. Vinokur, Physica **C254**, 63 (1995).